

Solid Electrolyte Interphase: Theory and Experiment

Birger Horstmann^{1,2}, Miriam Klein¹, Fabian Single^{1,2}, Erkmen Karaca^{1,2,3}, Norbert Wagner¹, K. Andreas Friedrich^{1,3}, Arnulf Latz

¹ German Aerospace Center, Stuttgart, Institute of Engineering Thermodynamics, Pfaffenwaldring 38-40, 70569 Stuttgart, Germany

² Helmholtz Institute Ulm, Helmholtzstraße 11, 89081 Ulm, Germany

³ University of Ulm, Albert-Einstein-Allee 11, 89081 Ulm, Germany

⁴ University of Stuttgart, Institute for Thermodynamics, Pfaffenwaldring 6, 70550 Stuttgart, Germany

Correspondence to birger.horstmann@dlr.de


Solid Electrolyte Interphase (SEI)

- Film formation on graphite anode
 - **Reduction product** of electrolyte
 - SEI slows down further electrolyte reduction
 - **Continued capacity fade** / Increase in impedance
- Challenges in Experiment and Theory
 - Complex **chemical composition**
 - SEI morphology: **porosity** / dual-layer structure
 - Limiting **transport process**

Experiment


- Graphite vs. Li/Li⁺
- Electrolyte: EC:DEC 3:7 wt.-%. 1 M LiPF₆
- Additive: Fluoroethylene carbonate (FEC), 2 wt.-%

Reaction of FEC¹:




Results:

Fourier Transform Infrared Spectroscopy



Cyclovoltammetry




Conclusion:

- VC is reduced earlier than the electrolyte (CV)
- A stable film on the anode side is formed
- Further electrolyte decomposition is prevented
- Homogeneity of SEI-layer verified with space-resolved FTIR

Modeling Scheme


- 1D mesoscopic continuum model on electrode surface
 - Simple SEI chemistry: Li₂CO₃
 - Electrolyte conductivity in **nanopores** $\sigma^{eff} = \epsilon^{\beta} \sigma$ with large empirical Brugemann coefficient β




- **Specific Surface Area** of porous SEI

$$A(\epsilon) = \frac{6}{a_0} \epsilon (1 - \epsilon) - a_0 \epsilon \frac{\partial^2 \epsilon}{\partial x^2}$$

- Elementary cubes of size a_0^3 filled with probability ϵ_{SEI}



Simulation and Discussion



- Numerical stability
- Front width scales with a_0

- **Thickness** growth with \sqrt{t} due to electron conductivity ($\kappa \approx 10^{-12}$ S/m)

$$\frac{dL}{dt} = \frac{\Phi' \kappa (1 - \epsilon(L))^{1.5}}{2F} (1 - \epsilon(L)) V^M$$

- **Constant porosity** ϵ_* due to balance with electrolyte transport ($\beta \approx 10$)

$$\beta \epsilon_*^{\beta-1} \sigma \approx (1 - \epsilon_*)^{2.5} \kappa$$

Literature

- [1] S. S. Zhang, J. Power Sources 162 (2006) 1379.
- [2] E. Peled, in Lithium Batteries, J.P. Gabao, Editor, Academic Press, New York (1983) 29.
- [3] M.B. Pinson, M.Z. Bazant, J. Electrochem. Soc. 160, A243 (2012) 30.
- [4] P. Lu et al., J. Phys. Chem. C 118, 896 (2014).
- [5] P. Liu et al., J. Electrochem. Soc. 157, A499 (2010).